

- [3] H. Doshi, A. Ray, and I. Kothari, "Biosorption of cadmium by live and dead *Spirulina*: IR spectroscopic, kinetics, and SEM studies," *Curr. Microbiol.*, vol. 54, pp. 213–218, 2007.
<https://doi.org/10.1007/s00284-006-0340-y>
- [4] J. Godt, F. Scheidig, C. Grosse-Siestrup, V. Esche, P. Brandenburg, A. Reich, and D. A. Groneberg. "The toxicity of cadmium and resulting hazards for human health," *Journal of occupational medicine and toxicology*; vol. 1, 2006.
- [5] Q. Hurdebise, C. Tarayre, C. Fischer, G. Colinet, S. Hiligsmann, and F. Delvigne, "Determination of zinc, cadmium and lead bioavailability in contaminated soils at the single-cell level by a combination of whole-cell biosensors and flow cytometry," *Sensors*, vol. 15, no. 4, pp. 8981-8999, 2015.
<https://doi.org/10.3390/s150408981>
- [6] I. Vopálenká, L. Váchová, and Z. Palková, "New biosensor for detection of copper ions in water based on immobilized genetically modified yeast cells," *Biosensors and Bioelectronics*, vol. 72, pp. 160-167, 2015.
<https://doi.org/10.1016/j.bios.2015.05.006>
- [7] H. Strosnider, "Whole-cell bacterial biosensors and the detection of bioavailable arsenic," *US Environmental Protection Agency*, 2003.
- [8] G. M. Walker, "Metals in yeast fermentation processes," *Advances in applied microbiology*, vol. 54, pp. 197-230, 2004.
[https://doi.org/10.1016/S0065-2164\(04\)54008-X](https://doi.org/10.1016/S0065-2164(04)54008-X)
- [9] T. D. Kaiser, E. M. Pereira, K. R. Dos Santos, E. L. Maciel, R. P. Schuenck, A. P. Nunes. "Modification of the Congo red agar method to detect biofilm production by *Staphylococcus epidermidis*," *Diagn Microbiol Infect Dis.*, vol. 75, pp. 235-9, 2013.
<https://doi.org/10.1016/j.diagmicrobio.2012.11.014>
- [10] N. Ö. Akpolat, S. Elci, S. Atmaca, H. Akbayin, and K. Gül, "The effects of magnesium, calcium and EDTA on slime production by *Staphylococcus epidermidis* strains," *Folia microbiologica*, vol. 48, pp. 649, 2003.
<https://doi.org/10.1007/BF02993473>